

Rotarix™ liquid oral vaccine

ROTAVIRUS VACCINE


Only 2 oral doses



Introducing the **NEW** flexible dosing schedule for Rotarix™

Easy implementation into existing immunisation schedules.


Dosage:



2 oral doses:

- First oral dose: as early as 6 weeks of age
- Second oral dose: completed by 24 weeks of age
 - Minimum interval between doses: 4 weeks

How to administer the oral liquid formulation



1. Remove the protective tip-cap from the oral applicator.



2. Administer the entire content of the applicator orally (on the inside of the cheek). The child should be seated in a reclining position. **Administer orally. Do not inject.**



DO NOT INJECT • FOR ORAL USE ONLY

[S2] ROTARIX™ LIQUID ORAL VACCINE. Rotavirus vaccine. Oral suspension. 43/30.2/0290. 1 dose (1,5 ml) contains: Live attenuated human rotavirus RIX4414 strain not less than 10^{6.0} CCID₅₀. **APPLICANT:** GlaxoSmithKline South Africa (Pty) Ltd. (Co. Reg. No. 1948/030135/07). 57 Sloane Street, Bryanston, 2021. Tel: +27 11 745 6000. Fax: +27 11 745 7000. For full prescribing information refer to the package insert approved by the medicines regulatory authority. Marketed by Aspen Pharmacare, Building 12, Healthcare Park, Woodlands Drive, Woodmead, 2191. www.aspenpharma.com. Medical Information Hotline 0800 118 088. ROT/0810/142 A1182 08/10



Healthcare. We Care.

